

Association Between Periodontal Pathogens and Severity of Periodontal Diseases Among Adolescents in Kunming City: China

Jie Xu, D.D.S., M.Sc., Ph.D.^{1,2,3}, Nuntiya Pahumunto, B.Sc., M.Sc., Ph.D.^{1,2},
Supawadee Naorungroj, D.D.S., M.Sc., Ph.D.^{1,4}, Rawee Teanpaisan, B.Sc., M.Sc., Ph.D.¹

¹Research Center of Excellence for Oral Health, Faculty of Dentistry, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand.

²Department of Oral Diagnostic Sciences, Faculty of Dentistry, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand.

³Department of Periodontology, the Affiliated Stomatology Hospital of Kunming Medical University, Yunnan, China.

⁴Department of Conservative Dentistry, Faculty of Dentistry, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand.

Received 18 January 2023 • Revised 14 June 2023 • Accepted 25 August 2023 • Published online 24 January 2024

Abstract:

Objective: To investigate the association between periodontal pathogens and periodontal status among adolescents in Kunming City.

Material and Methods: A total of 560 adolescents from five high schools within five urban districts in Kunming participated in this study. Clinical periodontal examination was assessed, and the levels of *Porphyromonas gingivalis* (*P. gingivalis*), *Aggregatibacter actinomycetemcomitans* (*A. actinomycetemcomitans*), *Prevotella intermedia* (*P. intermedia*), and *Fusobacterium nucleatum* (*F. nucleatum*) in saliva samples were examined using a real-time polymerase chain reaction (PCR). The relationship between periodontal pathogens and the severity of periodontal disease was investigated using multivariate logistic regression analysis.

Results: The presence of *P. gingivalis*, *A. actinomycetemcomitans*, and *P. intermedia* was positively correlated with the extent of probing depth, clinical attachment level (CAL), and bleeding on probing. *P. gingivalis* and *A. actinomycetemcomitans* were detected at approximately 55–58% among adolescents with healthy periodontal status. *P. gingivalis* and *A. actinomycetemcomitans* were detected in all adolescents diagnosed with periodontitis (stage I or II), which was an increased number compared to healthy adolescents. It was found that *P. gingivalis* (odds ratio (OR)=7.03, 95% confidence interval (CI) 3.77–13.11) and *A. actinomycetemcomitans* (OR=5.37, 95% CI 2.73–10.57) were strongly associated with stage II periodontitis. *F. nucleatum* had no significant relationship with the occurrence of gingivitis and periodontitis.

Conclusion: *P. gingivalis*, *A. actinomycetemcomitans*, and *P. intermedia* are important microbiological risk factors for periodontal diseases in adolescents.

Keywords: periodontal diseases, periodontal pathogens, *P. gingivalis*, risk factors

Contact: Rawee Teanpaisan, Ph.D.
Research Center of Excellence for Oral Health, Faculty of Dentistry,
Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand.
E-mail: rawee.t@psu.ac.th

J Health Sci Med Res 2024;42(3):e20241029
doi: 10.31584/jhsmr.20241029
www.jhsmr.org

© 2024 JHSMR. Hosted by Prince of Songkla University. All rights reserved.
This is an open access article under the CC BY-NC-ND license
(<http://www.jhsmr.org/index.php/jhsmr/about/editorialPolicies#openAccessPolicy>).

Introduction

Periodontal diseases are a group of diseases; including gingivitis and periodontitis, that are related to the inflammatory process. They are prevalent in many countries, and their prevalence and severity increase with age¹. Clinically, gingivitis often precedes periodontitis, with the detection of periodontal attachment loss^{2,3}. An epidemiological study reported that periodontal diseases could be found starting at a young age, 5–14 years old, with a high prevalence of gingivitis (84.4%)⁴. Another study showed that 31.3% of adolescents aged 12–15 in Shandong, China have gingivitis⁵. This recent study in Kunming, China also reported a high prevalence of gingivitis (80.4%) and periodontitis (7.3%) among adolescents aged 17–19⁶.

Childhood and adolescence are crucial periods for the development of periodontitis; therefore, it is crucial to comprehend the colonization and attachment of periodontal pathogens and their association with disease development. Many studies have identified risk factors for periodontal diseases; including an increase in the amount of certain periodontal microbiota^{7–11}. *Porphyromonas gingivalis* (*P. gingivalis*), *Aggregatibacter actinomycetemcomitans* (*A. actinomycetemcomitans*), *Prevotella intermedia* (*P. intermedia*), and *Fusobacterium nucleatum* (*F. nucleatum*) have been reported to play an important role in periodontal diseases occurrence and progression^{8,12}. However, the information relating to periodontopathogens among the Chinese population, particularly among adolescents, is scarce. Therefore, this study aimed to investigate the association between periodontal pathogens; including *P. gingivalis*, *A. actinomycetemcomitans*, *P. intermedia*, and *F. nucleatum*, and the periodontal status among adolescents in Kunming City.

Material and Methods

Study setting and participants

This was the second phase of a cross-sectional study regarding the prevalence and associated factors of dental caries and periodontal diseases among adolescents (aged 17–19) in Kunming High School. It was carried out from the academic year of July 2019 to November 2020. The study setting and participant recruitment have been published in our previous report¹², which has been approved by the Ethics Committee of Prince of Songkla University (EC6208–033) and Kunming Medical University (KY2020MEC019).

In brief, one public school from each of the five urban districts in Kunming was selected. High school students having lived in Kunming for more than a year were invited, and a total of 560 adolescents (17.3±0.5 years old) with an equal gender ratio gave consent to participate in this study. A self-administered questionnaire was then used in the survey to collect sociodemographic characteristics, health status, and oral health behaviors related to dental caries and periodontal diseases. Unstimulated whole saliva samples were collected before the periodontal examination.

Clinical periodontal examination

A full mouth clinical periodontal examination of all participants was carried out using a manual rigid periodontal probe (PCPUNC15, Hu Friedy, Chicago, IL, USA). Two calibrated examiners (Kappa value >0.8) assessed bleeding on probing (BOP), the probing depth (PD), and the clinical attachment level (CAL) at six sites per tooth (mesio-buccal, mid-buccal, disto-buccal, mesio-lingual, mid-lingual and disto-lingual). Clinical periodontal status was presented according to the 2018 AAP/EEP classification of periodontal disease into healthy, localized gingivitis, generalized gingivitis, stage I and stage II periodontitis¹³.

Following the clinical examination, the children were informed of their oral health status. Anyone who had gingivitis or periodontitis was advised to see a dentist.

Determination of periodontal pathogens using quantitative real-time polymerase chain reaction (PCR) assay

The unstimulated saliva from individual subjects was collected, transferred into a sterile container, and kept at -80°C until used for Deoxyribonucleic acid (DNA) extraction. Then, the saliva samples were extracted using the PureDirex Genomic DNA Isolation Kit (Bio-Helix Co., Ltd., Keelung, Taiwan); according to the manufacturer's instructions. The number of periodontal pathogens; including *A. actinomycetemcomitans*, *P. gingivalis*, *F. nucleatum*, *P. intermedia*, were determined using a 16S rRNA-based real-time polymerase chain reaction with specific primers (Table 1), and sterile deionized water was used as control.

The real-time PCR was performed in a 20- μL aliquot of real-time PCR master mix; including 5 μL of a purified DNA template, 5 pM of forward and reverse specific primers (Table 1), nuclease-free water, and 10 μL of 2x SensiFAST SYBR[®] No-ROX Mix (Bioline Reagent Ltd., California, USA), for PCR amplification. The real-time PCR conditions were as follows: denaturation at 95°C for 10 min and 50°C for 2 min, followed by 40 cycles at 95°C for 20s, 60°C for 20s, and 72°C for 25s. All samples were run in duplicate by a QuantStudio3 & QuantStudio5[™] Flex Real-Time PCR System (Thermo Fisher Scientific, USA). The number of targeting bacteria (*A. actinomycetemcomitans*, *P. gingivalis*, *F. nucleatum*, *P. intermedia*) in the saliva samples was calculated with a standard curve. For standard curve construction, bacterial cells were diluted in two-fold dilution and were aliquot into two tubes. The first aliquot counted the bacterial number using the plate count method (10^1 – 10^9 CFU/mL) and the second aliquot extracted DNA to measure the Ct value using QuantStudio3 & QuantStudio5[™] Flex Real-Time PCR System. A linear standard curve was plotted for each bacterial species from log CFU/mL against the corresponding Ct ($R^2 > 0.99$). The bacterial number was reported as log CFU/mL¹⁴.

Statistical methods

For overall analyses, the normal distribution and homogeneity of variance of continuous variables were examined. The median values of all log-transformed periodontal pathogen counts were calculated. As the distribution of the four periodontal pathogens' numbers was not normal, Spearman's rank correlation coefficient (r_s) was used to evaluate the correlation between the number of periodontal pathogens and the extent of periodontal clinical parameters (probing depth (PD), clinical attachment level (CAL), and bleeding on probing (BOP)). The bacteria detection frequency was estimated by dividing the number of subjects that tested positive for a specific bacterium, using real-time PCR by the total number of subjects. In addition, the correlation of the number of periodontal pathogens with different periodontal diseases statuses was analyzed by one-way analysis of variance (ANOVA) or nonparametric tests (Kruskal-Wallis test). Post-hoc pairwise comparisons with Bonferroni adjustment were conducted to determine significant differences between the periodontal diagnostic groups. Multivariate logistic regression analysis was used to investigate the relationship between periodontal pathogens and the severity of periodontal disease. All p -value < 0.05 were considered statistically significant.

Results

Spearman's rank correlation coefficients revealed a statistically significant positive correlation between the number of *P. gingivalis*, *A. actinomycetemcomitans*, and *P. intermedia* and three clinical periodontal parameters (BOP, PD, and CAL), indicating the level of gingival inflammation and periodontal destruction, as well as between the number of *F. nucleatum* and CAL. However, the correlations were weak ($r_s = 0.10$ – 0.36) (Table 2).

The prevalence and number of targeted periodontal pathogens in the saliva samples of 560 adolescents with different periodontal statuses, are demonstrated in

Table 3. Generally, the prevalence of all target periodontal pathogens; except *A. actinomycetemcomitans*, was relatively high among the Chinese adolescents; being 94.1%, 100%, 100%, and 59.6% for *P. gingivalis*, *P. intermedia*, *F. nucleatum*, and *A. actinomycetemcomitans*, respectively. It was found that the detection frequency and number of *P. gingivalis* and *A. actinomycetemcomitans* increased with the severity of periodontal status. They were substantially higher in adolescents with periodontitis and generalized gingivitis compared to those who were healthy or had localized gingivitis. In contrast, *P. intermedia* and *F. nucleatum* could be detected in all adolescents (100%); they were found to be higher in number than *P. gingivalis* and *A. actinomycetemcomitans* (Table 3).

In the multivariate logistic regression analysis for the association between periodontal pathogens and severity of periodontitis (Table 4), it was found that *P. gingivalis* was significantly associated with the occurrence of gingivitis (odds ratio (OR)=3.38, 95% confidence interval (CI) 2.40–4.75 for localized gingivitis, and OR=4.50, 95% CI 3.03–6.68 for generalized gingivitis and periodontitis (OR=3.20, 95% CI 1.75–5.84 for stage I, and OR=7.03, 95% CI 3.77–13.11 for stage II). An increased number of *A. actinomycetemcomitans* was associated with the increased odds of having generalized gingivitis (OR=3.32, 95% CI 1.91–5.77) and periodontitis (stage I and II). The odds ratio varied between 5.83–5.37. *P. intermedia* was associated only with the occurrence of periodontitis, with an odds ratio of 2.84–2.44. However, *F. nucleatum* had no significant relationship with the occurrence of gingivitis and periodontitis.

Table 1 Bacterial species and the real-time PCR primers used in this study

| Species | | Sequence (5'-3') | Product size |
|---------------------------------|----------------|-----------------------------|--------------|
| <i>P. gingivalis</i> | Forward primer | TGCAACTTGCCTTACAGAGGG | 344 bp |
| | Reverse primer | ACTCGTATCGCCCGTTATTC | |
| <i>A. actinomycetemcomitans</i> | Forward primer | GAACCTTACCTACTCTTGACATCCGAA | 80 bp |
| | Reverse primer | TGCAGCACCTGTCTCAAAGC | |
| <i>P. intermedia</i> | Forward primer | CCACATATGGCATCTGACGTG | 232 bp |
| | Reverse primer | CACGCTACTTGGCTGGTTCA | |
| <i>F. nucleatum</i> | Forward primer | GGATTTATTGGGCGTAAAGC | 163 bp |
| | Reverse primer | GGCATTCTACAAATATCTACGAA | |

PCR=polymerase chain reaction

Table 2 Correlation between periodontal pathogens and clinical parameters

| Clinical parameters | Spearman's rho (p-value) | | | |
|---------------------|--------------------------|---------------------------------|----------------------|---------------------|
| | <i>P. gingivalis</i> | <i>A. actinomycetemcomitans</i> | <i>P. intermedia</i> | <i>F. nucleatum</i> |
| PD | 0.20 (<0.001) | 0.24 (<0.001) | 0.14 (0.001) | 0.08 (0.063) |
| CAL | 0.20 (<0.001) | 0.30 (<0.001) | 0.20 (<0.001) | 0.10 (0.024) |
| BOP | 0.36 (<0.001) | 0.36 (<0.001) | 0.13 (0.003) | 0.02 (0.691) |

PD=probing depth, CAL=clinical attachment level, BOP=bleeding on probing

Table 3 Prevalence and number of periodontal pathogens among 560 adolescents having different periodontal statuses

| Periodontal pathogens | Total n=560 | Periodontal status | | | | |
|---------------------------------|----------------|--------------------|--------------------|----------------------|-----------------|------------------|
| | | Healthy n=69 | Gingivitis | | Periodontitis | |
| | | | Localized n=295 | Generalized n=155 | Stage I n=24 | Stage II n=17 |
| <i>P. gingivalis</i> | | | | | | |
| Frequency of detection, % | 94.10 | 55.10 | 99.30 | 100.00 | 100.00 | 100.00 |
| Log CFU/ml, Median | 2.04 | 0.36 | 1.89 | 2.45 | 2.34 | 2.85 |
| (min, max) | (0.00, 7.17) | (0.00, 0.27)*** | (0.00, 5.59)** | (0.07, 7.17)* | (0.04, 4.25)*** | (1.35, 4.13)* |
| <i>A. actinomycetemcomitans</i> | | | | | | |
| Frequency of detection, % | 59.60 | 58.00 | 46.80 | 74.20 | 100.00 | 100.00 |
| Log CFU/ml, Median | 0.40 | 0.13 | 0.00 | 1.12 | 2.24 | 2.01 |
| (min, max) | (0.00, 9.00) | (0.00, 9.00)** | (0.00, 3.07)** | (0.00, 3.16)* | (0.52, 8.09)* | (0.11, 3.24)* |
| <i>P. intermedia</i> | | | | | | |
| Frequency of detection, % | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 |
| Log CFU/ml, Median | 4.79 | 4.67 | 4.77 | 4.72 | 5.87 | 5.26 |
| (min, max) | (0.84, 7.19) | (0.91, 6.40)** | (0.84, 7.12)** | (1.40, 7.12)** | (3.10, 7.19)* | (2.58, 6.76)*** |
| <i>F. nucleatum</i> | | | | | | |
| Frequency of detection, % | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 | 100.00 |
| Log CFU/ml, Median | 5.32 | 5.36 | 5.33 | 5.24 | 5.73 | 5.72 |
| (min, max) | (0.12, 7.10) | (0.63, 6.47)* | (0.13, 7.10)* | (0.24, 6.66)* | (2.75, 6.65)* | (0.12, 6.47)* |

***, ** indicated the statistical difference between groups. One-way analysis of variance (ANOVA) was performed on variables with homogeneity of variance (*P. gingivalis*, *P. intermedia*, and *F. nucleatum*), and non-parametric test analysis (Kruskal-Wallis test) was performed on variables with unequal variance (*A. actinomycetemcomitans*). Post-hoc pairwise comparisons with Bonferroni adjustment were conducted to determine significant differences between the periodontal diagnostic groups.

Table 4 Multivariate logistic regression analysis of periodontal pathogens and different periodontal statuses (n=560)

| Periodontal pathogens | Localized gingivitis n=295 | | Generalized gingivitis n=155 | | Stage I periodontitis n=24 | | Stage II periodontitis n=17 | |
|---------------------------------|-------------------------------|---------|---------------------------------|---------|-------------------------------|---------|--------------------------------|---------|
| | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value | OR (95% CI) | p-value |
| <i>P. gingivalis</i> | 3.38 | <0.001 | 4.50 | <0.001 | 3.20 | <0.001 | 7.03 | <0.001 |
| | (2.40, 4.75) | | (3.03, 6.68) | | (1.75, 5.84) | | (3.77, 13.11) | |
| <i>A. actinomycetemcomitans</i> | 1.26 | 0.385 | 3.32 | <0.001 | 5.83 | <0.001 | 5.37 | <0.001 |
| | (0.75, 2.12) | | (1.91, 5.77) | | (3.07, 11.07) | | (2.73, 10.57) | |
| <i>P. intermedia</i> | 1.00 | 0.994 | 1.15 | 0.428 | 2.84 | 0.002 | 2.44 | 0.010 |
| | (0.74, 1.36) | | (0.81, 1.63) | | (1.45, 5.56) | | (1.24, 4.82) | |
| <i>F. nucleatum</i> | 1.02 | 0.942 | 1.23 | 0.404 | 1.08 | 0.834 | 0.73 | 0.297 |
| | (0.66-1.55) | | (0.76, 2.00) | | (0.52, 2.23) | | (0.40, 1.33) | |

The reference category is "Healthy" (n=69).
OR=odds ratio, 95% (CI)=95% confidence intervals

Discussion

Even though there are many studies investigating microbial species associated with periodontal diseases, most studies have been performed in adults with chronic periodontitis. This research examined the rate and number of periodontal pathogens in relation to the 2018 classification of periodontal diseases among Chinese adolescents. According to this study, the prevalence of gingivitis was 80.4% and periodontitis was 7.3%. The presence of *P. gingivalis* and *A. actinomycetemcomitans* is an important microbiological risk factor for periodontitis in adolescents.

Gingivitis and periodontitis are a result of gradually inflammatory processes, and certain bacteria play an important role in its initiation. In spite of a thousand species in the oral cavity, only some species; for example, *P. gingivalis* and *A. actinomycetemcomitans*, exhibited the potential to cause periodontal tissue destruction¹⁵. The main periodontal pathogens in this study included *P. gingivalis*, *A. actinomycetemcomitans*, *P. intermedia* and *F. nucleatum*. Overall, the bacterial prevalence obtained showed that most target periodontal pathogens except *A. actinomycetemcomitans*, were very high in prevalence among the Chinese adolescents; the higher prevalence of *P. gingivalis* and *A. actinomycetemcomitans* were found in the diseased stage (Periodontitis stage I and II) compared to the healthy. Furthermore, an increased severity of periodontitis was associated with an increase in *P. gingivalis* and *A. actinomycetemcomitans* levels (log CFU/mL). These results were very similar to a previous report;¹⁵ however, some discrepancies were noted in detail. In a study among 255 Indonesian adolescents with untreated periodontal disease, 57% had *A. actinomycetemcomitans*, 87% had *P. gingivalis*, and 89% had *P. intermedia*. The overall prevalence profile of periodontal pathogens was similar to this study. Furthermore, in that study, at site level *P. gingivalis* was more prevalent in sites with attachment

loss, which is consistent with this study's findings; in that *P. gingivalis* was more prevalent in subjects with stage I to stage II periodontitis. However, there was no significant association between the clinical periodontal parameters and the prevalence of microorganisms at the subject level in Indonesian adolescents¹⁶. Another study of periodontal pathogenic bacteria among high school children aged 15–18 years old in Saudi Arabia reported a relatively low prevalence of *A. actinomycetemcomitans* (21.7%), *P. gingivalis* (21.3%), *P. intermedia* (12.3%), and red complex bacteria (2.9%); however, there was no significance related to any clinical characters¹⁷.

Previous studies of periodontal microbiota and the clinical periodontal status among Thai and Chinese adults showed that the 'red complex' bacteria (*P. gingivalis*, *Tannerella forsythia*, and *Treponema denticola*) revealed a strong association between both pocketing and the extent of attachment loss. In addition, *P. gingivalis* was the one species that conferred the highest odds ratio^{18,19}. This is in agreement with a number of previous studies that reported *P. gingivalis* prevalence and levels significantly increased in adults with gingivitis and periodontitis^{20–23}. A study carried out among 116 Chinese with chronic periodontitis aged 20–75 years old reported that *A. actinomycetemcomitans* was more frequently detected in diseased sites than in periodontally healthy sites; its higher prevalence was more associated with severe sites than moderate and mild sites²⁴.

This study examined the association of periodontal pathogens in saliva and periodontal status among younger aged subjects (17–19 years old) using a 16S RNA real-time PCR. It should be stated that saliva can be used as a method for the detection of periodontal pathogens, which is in agreement with previous reports^{25,26}. Saliva sampling is non-invasive, painless, and easy to perform. It has been found that the presence and number of periodontal pathogens in saliva are correlated with those in dental

plaque²⁷. Results in this study showed that there was a strong, positive correlation between the levels of individual periodontal pathogens in saliva. The quantitative analysis of the correlation between periodontal pathogens and clinical periodontal status revealed that the increased levels of *P. gingivalis* and *A. actinomycetemcomitans* increased the severity of clinically periodontal diseases. Odds ratio analysis, according to the high levels of both those species, revealed a strong association with clinical periodontal status, and it became stronger in subjects with a higher degree of periodontitis. This suggests that *P. gingivalis* and *A. actinomycetemcomitans* may be suitable variables for the risk/causative role of the salivary microbiota in adolescents with periodontitis. The prevalence and level of *F. nucleatum* were high in all subjects; however, there was no association with any clinical parameters and status. As known, *F. nucleatum* acts as a bridge organism between early and late colonizers of dental plaque formation²⁸ although its specific role in periodontal diseases may need clarification. Based on the findings of this study, it is reasonable to monitor the levels of certain salivary microbiota related to periodontal conditions at an early age, because it can provide timely preventive and/or therapeutic measures for periodontal diseases at an early stage. In this study, *P. gingivalis* and *A. actinomycetemcomitans* could be found in healthy children; however, they were in low levels of frequency and numbers compared to children with diseases. Due to the high sensitivity of the real-time PCR used, a low number of microorganisms could be detected. However, this may not necessarily result in the onset of periodontal diseases since the occurrence of diseases involves multifactors such as immunity, personal habits, oral hygiene and so forth. However, those children should be monitored.

Within the limitations of this cross-sectional study, it is difficult to explain any differences that may result from different factors, for example, ethnicity, age, a technique

used for microbial detection, clinical parameters and the classification system. A longitudinal study on the role of periodontal pathogens in an adolescent population will provide more understanding of the development of periodontitis and could provide new insights for periodontitis monitoring and prevention strategies.

Conclusion

In conclusion, levels of *P. gingivalis*, *A. actinomycetemcomitans*, and *P. intermedia*, but not *F. nucleatum*, were significantly correlated with clinical periodontal conditions. *P. gingivalis*, *A. actinomycetemcomitans*, and *P. intermedia* are significant risk factors for the development of periodontal disorders in adolescents based on their higher odds ratios.

Ethical approval of research

The research protocol was approved by the Ethics Committee of Research Prince of Songkla University (EC6208-033) and Kunming Medical University (KY2020MEC019)

Acknowledgement

We are very grateful for all school for their contributions in this study.

Funding sources

This work was supported by a grant from the Prince of Songkla University (Graduate Studies) for the Ph.D. study of Jie Xu, and from the School of Stomatology, Kunming Medical University.

Conflicts of interest

The authors have no conflicts of interest.

References

- American Academy of Periodontology–Research, Science and therapy Committee. Periodontal diseases of children and adolescents. *Pediatr Dent* 2008;30:240–7.
- Loe H, Morrison E. Periodontal health and disease in young people: screening for priority care. *Int Dent J* 1986;36:162–7.
- Page RC, Kornman KS. The pathogenesis of human periodontitis: an introduction. *Periodontol 2000* 1997;14:9–11.
- Dhar V, Jain A, Van Dyke TE, Kohli A. Prevalence of gingival diseases, malocclusion and fluorosis in school-going children of rural areas in Udaipur district. *J Indian Soc Pedod Prev Dent* 2007;25:103–5.
- Zhang M, Lan J, Zhang T, Sun W, Liu P, Wang Z. Oral health and caries/gingivitis-associated factors of adolescents aged 12–15 in Shandong province, China: a cross-sectional oral health survey. *BMC Oral Health* 2021;21:1–8.
- Xu J, Naorungroj S, Pahumunto N, Teanpaisan R. Prevalence and associated factors of caries and periodontal diseases among adolescents in Kunming: a cross-sectional study. *J Health Sci Med Res* 2023;41:e2022903
- Albandar JM. Global risk factors and risk indicators for periodontal diseases. *Periodontol 2000* 2002;29:177–206.
- Socransky SS, Haffajee AD, Cugini MA, Smith C, Kent Jr RL. Microbial complexes in subgingival plaque. *J Clin Periodontol* 1998;25:134–44.
- Haffajee AD, Socransky SS. Microbial etiological agents of destructive periodontal diseases. *Periodontol 2000* 1994;5: 78–111.
- Ledder RG, Gilbert P, Huws SA, Aarons L, Ashley MP, Hull PS, et al. Molecular analysis of the subgingival microbiota in health and disease. *Appl Environ Microbiol* 2007;73:516–23.
- Parahitiyawa NB, Scully C, Leung WK, Yam WC, Jin LJ, Samaranyake LP. Exploring the oral bacterial flora: current status and future directions. *Oral Dis* 2010;16:136–45.
- Hajishengallis G, Darveau RP, Curtis MA. The keystone-pathogen hypothesis. *Nat Rev Microbiol* 2012;10:717–25.
- Papapanou PN, Sanz M, Buduneli N, Dietrich T, Feres M, Fine DH, et al. Periodontitis: Consensus report of workgroup 2 of the 2017 world workshop on the classification of periodontal and peri-implant diseases and conditions. *J Periodontol* 2018;89:173–82.
- Manmontri C, Nirunsittirat A, Piwat S, Wattanarat O, Pahumunto N, Makeudom A, et al. Reduction of *Streptococcus mutans* by probiotic milk: a multicenter randomized controlled trial. *Clin Oral Investig* 2020;24:2363–74.
- Slots J, Ting M. *Actinobacillus actinomycetemcomitans* and *Porphyromonas gingivalis* in human periodontal disease: occurrence and treatment. *Periodontol 2000* 1999;20:82–121.
- Timmerman MF, Van der Weijden GA, Armand S, Abbas F, Winkel EG, Van Winkelhoff AJ, et al. Untreated periodontal disease in Indonesian adolescents. Clinical and microbiological baseline data. *J Clin Periodontol* 1998;25:215–24.
- Alghamdi AS, Almarghani AA. Periodontal pathogenic bacteria among high school children in Saudi Arabia. *Ann Saudi Med* 2019;39:244–50.
- Papapanou PN, Baelum V, Luan WM, Madianos PN, Chen X, Fejerskov O, et al. Subgingival microbiota in adult Chinese: prevalence and relation to periodontal disease progression. *J Periodontol* 1997;68:651–66.
- Papapanou PN, Teanpaisan R, Obiechina NS, Pithpornchaiyakul W, Pongpaisal S, Pisuthanakan S, et al. Periodontal microbiota and clinical periodontal status in a rural sample in southern Thailand. *Eur J Oral Sci* 2002;110:345–52.
- Boutaga K, van Winkelhoff AJ, Vandenbroucke-Grauls CM, Savelkoul PHM. The additional value of real-time PCR in the quantitative detection of periodontal pathogens. *J Clin Periodontol* 2006;33:427–33.
- Masunaga H, Tsutae W, Oh H, Shinozuka N, Kishimoto N, Ogata Y. Use of quantitative PCR to evaluate methods of bacteria sampling in periodontal patients. *J Oral Sci* 2010;52:615–21.
- Lau L, Sanz M, Herrera D, Morillo JM, Martín C, Silva A. Quantitative real-time polymerase chain reaction versus culture: a comparison between two methods for the detection and quantification of *Actinobacillus actinomycetemcomitans*, *Porphyromonas gingivalis* and *Tannerella forsythensis* in subgingival plaque samples. *J Clin Periodontol* 2004;31:1061–9.
- Kuboniwa M, Amano A, Kimura KR, Sekine S, Kato S, Yamamoto Y, et al. Quantitative detection of periodontal pathogens using real-time polymerase chain reaction with TaqMan probes. *Oral Microbiol Immunol* 2004;19:168–76.

24. Meng S, Zhao L, Yang H, Wu Y, Ouyang Y. Prevalence of *Actinobacillus actinomycetemcomitans* in Chinese chronic periodontitis patients and periodontally healthy adults. *Quintessence Int* 2009;40:53–60.
25. Umeda M, Contreras A, Chen C, Bakker I, Sots J. The utility of whole saliva to detect the oral presence of periodontopathic bacteria. *J Periodontol* 1998;69:828–33.
26. Boutaga K, Savelkoul PH, Winkel EG, van Winkelhoff AJ. Comparison of subgingival bacterial sampling with oral lavage for detection and quantification of periodontal pathogens by real-time polymerase chain reaction. *J Periodontol* 2007;78:79–86.
27. He J, Huang W, Pan Z, Cui H, Qi G, Zhou X, et al. Quantitative analysis of microbiota in saliva, supragingival, and subgingival plaque of Chinese adults with chronic periodontitis. *Clin Oral Investig* 2012;16:1579–88.
28. Bolstad AI, Jensen HB, Bakken V. Taxonomy, biology, and periodontal aspects of *Fusobacterium nucleatum*. *Clin Microbiol Rev* 1996;9:55–71.