Adnexal Torsion During Controlled Ovarian Stimulation: A Case Report

Ekasak Thiangphak, M.D., Phawat Matemanosak, M.D., Suwadee Wirotkarun, B.Sc.

Department of Obstetrics and Gynaecology, Faculty of Medicine, Prince of Songkla University, Hat Yai, Songkha 90110, Thailand. Received 12 April 2024 • Revised 29 July 2024 • Accepted 19 August 2024 • Published online 24 February 2025

Abstract:

A 34-year-old nulliparous woman with polycystic ovarian syndrome presented with mild discomfort in her right abdomen three hours before her scheduled retrieval. She underwent a transvaginal oocyte retrieval. Oocytes were obtained from the ovary; however, the right ovary was inaccessible as it was located above the uterine fundus. Consequently, the decision was made to abandon retrieval from the right ovary. After the procedure, the pain continued, and a physical examination indicated signs of peritonitis. Right ovarian torsion was suspected, leading to laparoscopic detorsion with transabdominal oocyte retrieval from the untwisted ovary. One mature oocyte was retrieved from the torsion-affected ovary, resulting in an embryonic arrest at the morula stage. Additionally, we reviewed the literature on the outcomes of in vitro fertilization cycles in women with ovarian torsion before oocyte retrieval.

Keywords: in vitro fertilization, ovarian stimulation, ovarian torsion

Contact: Phawat Matemanosak, M.D. Department of Obstetrics and Gynaecology, Faculty of Medicine, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand. E-mail: phawattm@gmail.com J Health Sci Med Res 2025;43(5):e20251155 doi: 10.31584/jhsmr.20251155 www.jhsmr.org

© 2025 JHSMR. Hosted by Prince of Songkla University. All rights reserved. This is an open access article under the CC BY-NC-ND license (http://www.jhsmr.org/index.php/jhsmr/about/editorialPolicies#openAccessPolicy).

Introduction

Ovarian torsion during in-vitro fertilization (IVF) is a rare event, with an incidence of 0.024%–0.2%^{1,2}. Women with ovarian hyperstimulation syndrome (OHSS) and pregnancy after assisted reproduction are at a higher risk of torsion due to increased ovarian size². Delays in diagnosis and management can lead to compromised ovarian function and future fertility³. Therefore, ovarian torsion should be suspected and ruled out in women undergoing fertility treatment that present with sudden-onset severe abdominal pain.

Data on ovarian torsion occurring before oocyte retrieval regarding outcomes of surgery and IVF treatment are limited⁴⁻⁹. To our knowledge, this is the first case report of ovarian torsion during an IVF cycle in which laparoscopic detorsion was performed, followed immediately by oocyte retrieval from the affected ovary within the same procedure. Despite a delay in the timing of the retrieval, a morula-stage embryo was successfully obtained.

Case report

This study was approved by the Institutional Review Board of the Faculty of Medicine at the Prince of Songkla University (approval number: REC. 66–272–12–1). Patient approval for publication was obtained. A 34-year-old nulliparous woman with a BMI of 28 kg/m² presented at the clinic with a 3-year history of primary infertility related to polycystic ovarian syndrome. She had history of irregular menstruation, with an interval of 2–3 months. Pelvic ultrasonography showed an antral follicle count of >20 bilaterally and an ovarian volume of 15 and 10 mL on the left and right sides, respectively, consistent with polycystic ovarian morphology. Other investigations and her medical history were unremarkable.

Choice of fertility treatment was offered, and she proceeded with her first IVF cycle, using progestin-primed ovarian stimulation protocol. Follitropin alfa (r-hFSH-alfa, GONAL-f[®]; Merck, KGaA, Darmstadt, Germany) at a daily dose of 150 IU was started on day 2 of progestin-induced withdrawal bleeding. A 10-mg dose of medroxyprogesterone acetate per day was administered to prevent premature luteinizing from cycle day 3 onwards. The peak estrogen level on the day of the trigger (cycle day 10) was 10,124 pg/mL, GnRH-agonist (Diphereline, France, Epson) upon which 0.2 mg was administered for final oocyte maturation. She complained of mild discomfort in the right side of her abdomen three hours before her planned retrieval. A differential diagnosis of premature ovulation and ovarian torsion combined with OHSS was determined. She underwent transvaginal oocyte retrieval, and an ultrasound performed during the procedure revealed an anteverted uterus, bilateral enlarged multicystic ovaries, and minimal free fluid in the cul-de-sac. The left ovary was in a cul-desac, and the right ovary was positioned above the uterine fundus. Color Doppler ultrasonography was inconclusive in diagnosing adnexal torsion. Eleven oocytes were retrieved from the left ovary. The transvaginal approach to aspirate oocytes from the right ovary was attempted; however, only two follicles were aspirated due to the abnormal position of the ovary. Thus, oocyte retrieval from the remaining follicles of the right ovary was abandoned.

After oocyte retrieval, the pain persisted and she had severe abdominal pain on her right side, with increasing severity within the next hour. Her vital signs were as follows: pulse rate, 92 beats/min; body temperature, 36.8 °C, and blood pressure, 110/85 mmHg. Physical examination revealed moderate abdominal distension, guarding, and rebound tenderness around the lower right abdomen. Laboratory investigation revealed leukocytosis, with an elevated white blood cell count of 16,100 per microliter. Transabdominal ultrasound showed the left ovary measuring 7.6×6.5 cm in a cul-de-sac and the right ovary measuring 8.5×8.6 cm positioned high in the midline above the uterine fundus. Color Doppler ultrasonography was inconclusive in diagnosing ovarian torsion. However, right ovarian torsion was diagnosed through a combination of history and physical examination. Hence, an emergency laparoscopy for adnexal detorsion was performed 6 hours after symptom onset (3 hours after oocyte retrieval).

Laparoscopic approaches revealed a right enlarged ovarian cyst, measuring 8×8 cm, with an area of hemorrhage (Figure 1). The right adnexa had undergone 540° torsion around the utero-ovarian ligament. The twisted ovarian cyst was gently grasped with atraumatic forceps and detorsion was performed. Under laparoscopic vision a 17-gauge, single-lumen needle connected to an aspiration pump was used for transabdominal oocyte retrieval. The ovarian fluid was aspirated; however, only a minimal amount of follicular fluid was collected. At the end of the procedure, both adnexa were re-anatomized into a cul-de-sac.

Postoperatively her abdominal pain had improved, and as there were no postoperative complications, the patient was discharged 4 days after the operation. The follicular fluid from the two follicles of the right ovary was blood-stained and yellowish in color, and no oocytes were retrieved. One mature oocyte was retrieved from the torsion-affected ovary after detorsion was performed, which resulted in an embryonic arrest at the morula stage. Eleven oocytes were retrieved from the left ovary (9 MII and 2 GV), yielding three high-quality blastocysts; these were cryopreserved on day 5.

Discussion

The patient experienced ovarian torsion during ovarian stimulation after the trigger, and underwent transvaginal oocyte retrieval with subsequent laparoscopic detorsion in addition to transabdominal oocyte retrieval from a twisted ovary.

Adnexal torsion during fertility treatment is rare; however, it is a gynecological emergency requiring urgent surgical management to preserve ovarian function^{1,2}. Adnexal torsion can be diagnosed when taking into account the patient's history, physical examination, and imaging studies. Common presenting signs and symptoms include: sudden onset of severe abdominal pain, nausea, and vomiting. Color Doppler ultrasound findings such as enlarged



Figure 1 (a) Intraoperative findings demonstrating an enlarged edematous right ovary, with focal area of hemorrhage. (b) A single-lumen needle was used for transabdominal aspiration of the follicular fluid

or edematous ovaries, decreased or absent venous flow, follicular ring sign and whirlpool sign, are suggestive of torsion¹⁰. However, normal Doppler flow cannot reliably rule out adnexal torsion due to its low sensitivity. This limitation arises because arterial flow might still be preserved in cases with early presentation^{11,12}: as was observed in our case. Hence, adnexal torsion is a clinical diagnosis, and a high index of suspicion for torsion is essential.

The majority of reports of ovarian torsion during ovarian stimulation occurred after oocyte retrieval, especially after embryo transfer, and patients usually have OHSS^{2,13-15}. Data on ovarian torsion occurring before oocyte retrieval regarding outcomes of IVF treatment are limited⁴⁻⁹ (Table 1). Cases where torsion occurs before retrieval show a decrease in the number of oocytes retrieved as well as the fertilization rate of oocytes retrieved from a twisted ovary. Several mechanisms leading to decreased fertilization rates of oocytes from twisted ovaries have been proposed; including a decreased arterial blood supply to the ovary, leading to a reduced concentration of essential hormones in folliculogenesis; venous and lymphatic stasis contributing to dysfunctional folliculogenesis⁶, and ischemic injury to oocytes⁴.

The outcome of IVF, where torsion occurs before the planned retrieval and oocytes are retrieved prior to performing laparoscopic detorsion, is poorer than for those who undergo laparoscopic detorsion followed by oocyte retrieval. This aligns with our case. Stefanidis et al⁶. reported a case of a 33-year-old female who presented with abdominal pain one day before her scheduled retrieval. The patient proceeded directly to oocyte retrieval as planned. Twelve follicles were aspirated from the twisted ovary; however, no oocytes were obtained. The patient's symptoms worsened postoperatively, leading to a laparoscopy performed six hours after oocyte retrieval. Robson et al⁵. also reported a case of a 37-year-old female patient whose symptoms arose one hour before her planned retrieval. Oocyte retrieval was conducted to aspirate all follicles and reduce ovarian volume, which led to the resolution of the patient's symptoms. The fertilization rate from the affected side was lower than that of the contralateral side (86% vs. 33%). The patient's pain intensified, prompting a laparoscopy two hours later.

Conservative surgical management with detorsion under laparoscopy represents the best option among women with adnexal torsion that occurs before the time of planned retrieval. Detorsion should be performed before proceeding directly to oocyte retrieval. Proceeding directly to oocyte retrieval with the aim of reducing ovarian volume to obliviate the surgery was unsuccessful in two previous studies, resulting in the need for subsequent laparoscopic detorsion^{5,6}. Similar to our case, this approach necessitated subsequent surgery after ovum pick-up. Moreover, the outcome of IVF showed a decrease in the fertilization rate⁵ and empty follicle syndrome from the twisted ovary⁶. In contrast, one study reported success in reducing ovarian volume by aspiration when ovarian torsion was suspected two days before oocyte retrieval. This relieved the patient's pain and allowed for successful oocyte collection two days later. Notably, laparoscopy was not performed to confirm torsion in this instance¹⁴.

This case raises important points. First, ovarian torsion during IVF is a rare but serious complication. A high index of suspicion and early diagnosis are crucial for preserving ovarian function. Second, laparoscopic detorsion should be performed before proceeding directly to oocyte retrieval if torsion occurs prior to the scheduled retrieval time. Finally, in situations where diagnosis delays result in postponed oocyte retrieval, aspiration of any remaining follicles to reduce ovarian volume might lower the chance of re-torsion. Hence, the possibility of obtaining viable oocytes could still be attempted. Table 1 A summary of in vitro fertilization outcomes from published case reports regarding ovarian torsion before oocyte retrieval

Case report	Patient profile and	Onset of symptoms	Management	Cycle	outcome
	cause of infertility		•	Affected ovary	Unaffected ovary
Robson, et al. 2000	 - 37 year-old nulligravida - Male factor 	One hour before the planned retrieval	Oocyte retrieval followed by laparoscopic detorsion two hours after oocyte retrieval	 7 oocytes were retrieved 33% fertilization rate 	 12 oocytes were retrieved 86% fertilization rate 3 embryos were transferred (data on transfer outcome are unavailable)
Stefanidis, et al. 2002	 33-year-old nulligravida Male factor 	One day before the planned retrieval	Oocyte retrieval followed by laparoscopic detorsion six hours after oocyte retrieval	- 12 follicles were aspirated but no oocytes were retrieved	 18 oocytes were retrieved Transfer of 4 embryos; however pregnancy did not ensue
Smith, et al. 2010	 27-year-old nulligravida Husband carried a balanced translocation 	One day before the planned retrieval	Laparoscopic detorsion followed by oocyte retrieval on the next day	 10 oocytes were retrieved 4./10 (40%) were fertilized 3./4 (75%) developed into blastocysts All embryos were biopsied and revealed unbalanced translocations None were transferred 	 15 occytes were retrieved 14/15 (33%) were fertilized 9/14 (64%) developed into blastocysts All embryos were biopsied and revealed unbalanced translocations None were transferred
Inoue D. and Asada Υ. 2020	- 34-year-old nulligravida	Two days before oocyte retrieval	Ovarian volume reduction by aspiration two days before oocyte retrieval and successful oocyte collection two days later	 6 MII oocytes were retrieved 14.15 (93%) were fertilized 8 two-pronuclear (2PN) embryos were cryc the blastocyst stage. Two blastocysts were transferred, resulting 	 9 MII oocytes were retrieved opreserved, while 6 were cultured to in a successful pregnancy.
Naert, et al. 2023	 - 32-year-old nulligravida - Polycystic ovarian syndrome 	On cycle day 13 (one day before trigger)	Laparoscopic detorsion followed by trigger on the next day, and oocyte retrieval 36 hours after trigger	 3 mature oocytes were retrieved 2 blastocysts; one with high quality for cryopreservation 	 - 8 mature oocytes were retrieved - 5 blastocysts were cryopreserved on day 5 - 2 blastocysts were transferred resulting in live birth
Mortimer, et al. 2023	 - 34-year-old nulligravida - Polycystic ovarian syndrome 	12 hours before the planned retrieval	Laparoscopic detorsion followed by oocyte retrieval on the next day	 A total of 72 oocytes were retrieved, of wh oocytes. Out of the 70 MII oocytes, 36 were cryopreusing conventional IVF, resulting in a fertilius 16 blastocyst-stage embryos were cryopre. (The author did not specifically report the c 	ich 70 were metaphase II (MII) sserved, and 34 were inseminated zation rate of 79.4% (27 out of 34). served.
Our case	 - 34-year-old nulligravida - Polycystic ovarian syndrome 	Three hours before the planned retrieval	Oocyte retrieval followed by laparoscopic detorsion and second oocyte retrieval three hours later	 2 Follicles were aspirated in the first retrieval but no oocytes were retrieved 1 mature oocyte was retrieved from the second retrieval and developed into a morula-staged embryo 	 9 mature oocytes were retrieved 9/9 (100%) were fertilized 3/9 (33%) developed into blastocysts

Adnexal Torsion and Oocyte Retrieval

Conclusion

A high index of suspicion is required for diagnosing ovarian torsion occurring during IVF prior to ovum pickup; so that surgery can be promptly performed to improve the outcome of IVF cycles. However, if there is a delayed diagnosis, viable oocytes can still be obtained from the affected ovary after laparoscopic detorsion; despite potential delays in the timing of retrieval.

Acknowledgement

The authors would like to thank the patient.

Conflicts of interest

The authors report that there are no competing interests to declare.

References

- Bodri D, Guillén JJ, Polo A, Trullenque M, Esteve C, Coll O. Complications related to ovarian stimulation and oocyte retrieval in 4052 oocyte donor cycles. Reprod Biomed Online 2008;17:237–43.
- Spitzer D, Wirleitner B, Steiner H, Zech NH. Adnexal torsion in pregnancy after assisted reproduction – case study and review of the literature. Geburtsh Frauenh 2012;72:716–20.
- Krishnan S, Kaur H, Bali J, Rao K. Ovarian torsion in infertility management – Missing the diagnosis means losing the ovary: A high price to pay. J Hum Reprod Sci 2011;4:39–42.
- Smith LP, Oskowitz SP, Barrett B, Bayer SR. IVF and embryo development subsequent to ovarian torsion occurring during the resumption of meiosis. Reprod Biomed Online 2010;21:418–21.
- Robson S, Kerin JF. Acute adnexal torsion before oocyte retrieval in an in vitro fertilization cycle. Fertil Steril 2000;73:650–1.

- Stefanidis K, Grammatis M, Pappas K, Haidopoulos D, Chronopoulos K, Pantos K, et al. Empty follicle syndrome associated with ovarian torsion in an in vitro fertilization program. JSLS 2002;6:215–6.
- Inoue D, Asada Y. Successful oocyte retrieval after follicular fluid aspiration in suspicious of ovarian torsion. Cureus 2020;12:e12192.
- Naert MN, Baker S, Attaman J, Morris ME, Souter I, Fitz VW. Ovarian torsion during ovarian stimulation prior to trigger followed by successful oocyte retrieval: a case report. Case Rep Womens Health 2023;10:e00495.
- Mortimer RM, Gordon C, Srouji S. In vitro fertilization cycle outcomes for a patient undergoing oocyte retrieval following ovarian torsion. CRSLS 2023;10:e2022.00092.
- Chang-Patel EJ, Palacios-Helgeson LK, Gould CH. Adnexal torsion: a review of diagnosis and management strategies. Curr Opin Obstet Gynecol 2022;34:196–203.
- Nizar K, Deutsch M, Filmer S, Weizman B, Beloosesky R, Weiner Z. Doppler studies of the ovarian venous blood flow in the diagnosis of adnexal torsion. J Clin Ultrasound 2009;37:436– 9.
- Mashiach R, Melamed N, Gilad N, Ben-Shitrit G, Meizner I. Sonographic diagnosis of ovarian torsion: accuracy and predictive factors. J Ultrasound Med 2011;30:1205-10.
- Govaerts I, Devreker F, Delbaere A, Revelard P, Englert Y. Short-term medical complications of 1500 oocyte retrievals for in vitro fertilization and embryo transfer. Eur J Obstet Gynecol Reprod Biol 1998;77:239–43.
- Gorkemli H, Camus M, Clasen K. Adnexal torsion after gonadotrophin ovulation induction for IVF or ICSI and its conservative treatment. Arch Gynecol Obstet 2002;267:4–6.
- Zhu W, Li X, Chen X, Fu Z. Conservative management of adnexal torsion via transvaginal ultrasound guided ovarian cyst aspiration in patients with ovarian hyperstimulation. Fertil Steril 2008;89:229.e1–3.