

## Self-Care and Family Support among People with Type 2 Diabetes

Shashi Kandel, M.B.B.S.<sup>1</sup>, Wit Wichaidit, Ph.D.<sup>2</sup>

<sup>1</sup>Department of Health Services, Ministry of Health and Population, Government of Nepal, Kathmandu, Nepal.

<sup>2</sup>Epidemiology Unit, Faculty of Medicine, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand.

Received 26 March 2020 • Revised 12 May 2020 • Accepted 16 May 2020 • Published online 10 August 2020

### Abstract:

**Objective:** To assess the prevalence of self-care practices, and family support among people with type 2 diabetes.

**Material and Methods:** We conducted a cross-sectional study among outpatients for people with type 2 diabetes, at a teaching hospital in Kathmandu, Nepal. We modified the Summary of Diabetes Self-Care Activities, and Diabetes Family Behavior Checklist scales; to measure self-care practices and perceived family support, respectively, so as to better suit the Nepali context. We performed an exploratory factor analysis to identify independent, latent dimensions of family support.

**Results:** We recruited 411 patients (56.9% females), aged between 25 and 92. From these, we found high levels of adherence to daily intakes of the oral hypoglycemic agent (98.2%) and insulin (100.0%), daily physical activity (86.4%) and daily vegetable consumption (78.3%). Respondents commonly reported that they, and their family members ate meals together (98.1%). Family members also suggested things to help them take their medications on time (89.5%) and additionally helped them to decide on changes based on their glucose test results (84.2%). Exploratory factor analysis uncovered three behavior dimensions: nagging and arguing (“authoritarian”), praise and encouragement (“supportive”) and plan-making (“planning”).

**Conclusion:** We presented data on self-care as well as family support among people with type 2 diabetes, tailored to the local context. From the data a high level of medication adherence and daily physical activity was observed. “Eating together” was the most reported family support behavior. We identified three dimensions of family support behaviors, which are potentially helpful for future health promotion programs, and provides a methodological basis for future studies in other low and middle-income countries.

**Keywords:** family support, self-care practices, type 2 diabetes

**Contact:** Shashi Kandel, M.B.B.S.  
Department of Health Services, Ministry of Health and Population,  
Government of Nepal, Kathmandu, Nepal.  
E-mail: 6110320019@psu.ac.th

J Health Sci Med Res 2021;39(1):23-33  
doi: 10.31584/jhsmr.2020756  
www.jhsmr.org

© 2020 JHSMR. Hosting by Prince of Songkla University. All rights reserved.  
This is an open access article under the CC BY-NC-ND license  
(<http://www.jhsmr.org/index.php/jhsmr/about/editorialPolicies#openAccessPolicy>).

## Introduction

In 2017, there were 425 million adults, aged 20 to 70, who were suffering from diabetes globally; 90.0% of whom had type 2 diabetes mellitus (DM).<sup>1</sup> Control of blood glucose levels, also known as glycemic control, helps to prevent or delay DM-related microvascular and macrovascular complications.<sup>2</sup> Self-care practices that contribute to achieving glycemic control and reducing complications in people with type 2 diabetes include medication adherence, dietary behaviors, physical activity, self-monitoring of blood glucose levels and foot care.<sup>3-5</sup> However, adherence to self-care practices may vary by activity<sup>6</sup>, and self-care practices in each country vary in forms along with prevalence; based on each country's context.

Family support is the support provided by any family member to a person having a disability, which may include a child, an adult, or a parent. Many studies have shown that family support can improve self-care among diabetic patients.<sup>7-9</sup> However, the expression and function of family support may vary according to culture.<sup>10-12</sup> Culturally, nuanced care, strategies, and meaningful approaches are to be considered for better self-care management among people with type 2 diabetes. However, previous studies on family support have been conducted in high-income countries<sup>7-9</sup>, and the findings may not be generalizable to low- and middle-income countries.

Little information is available about diabetes self-care practices and family support among people with type 2 diabetes in Nepal; a low-income country in South Asia, undergoing an epidemiological transition. Such information can contribute to improvement in diabetes health promotion program designs and planning. This study; therefore, aimed to assess the prevalence of self-care practices and family support among people with type 2 diabetes in Nepal.

## Material and Methods

We conducted a hospital-based, cross-sectional study in Kathmandu, Nepal; from October to December, 2019, at Tribhuvan University Teaching Hospital's (TUTH) outpatient department, among people with type 2 diabetes. The TUTH's endocrinology outpatient department runs three days per week, and the general practice outpatient provides service six days per week. Both outpatient department receives, manage, and follow-up the people with type 2 diabetes.

We calculated the sample size based on an estimate of the proportion of participants having a high level of diabetes self-care behaviors. No previous estimate for this proportion could be found in the literature, so we assumed that 50.0% of people with type 2 diabetes had high adherence to self-care behaviors. Using a 95.0% level of confidence and a 5.0% margin of error, a sample size of 385 was deemed sufficient. We added a 10.0% non-response rate to the calculated sample size; hence, the final sample size was increased to 428 individuals.

After contacting the hospital administrator and explaining the objectives of the study, permission was given, and letters were sent to the heads of the Departments of Endocrinology and General Practice. Verbal permission to conduct the study along with access to the medical records was granted from each department. Using the medical records, we identified potential participants from the list of outpatients who visited the hospital; whom were expected to return for follow-up visits and diabetes management. Inclusion criteria included: (1) age 18 years or over; (2) attending the outpatient department; (3) previous diagnosis with type 2 DM for 3 months or longer at the time of the study, and (4) able to communicate unassisted in the Nepali language. We excluded patients who were dependent on other family members for daily living, those with cognitive

impairment or nervous system disability, those with serious comorbidity and those who lived alone. Data collectors approached potential participants, informed them about the study, and asked those who showed a willingness to participate for their written informed consent. After the participant read and signed the informed consent form, they were interviewed using a structured questionnaire.

Modified versions of the summary of diabetes self-care activities (SDSCA)<sup>13</sup> coupled with the diabetes family behavior checklist (DFBC)<sup>14</sup> were used to measure self-care practices and perceived family support, respectively. These instruments were first translated by an expert from English to Nepali, and then translated back to English by another expert; whom had not been exposed to the source documents. Discrepancies between the two English versions were used to identify potential translation issues in the Nepali version, which was then resolved by the investigators and the translators to obtain the final version.

#### **Measurement of diabetes self-care behaviors**

Self-care behaviors included five domains: (1) medication adherence; (2) physical activity; (3) dietary behaviors; (4) self-monitoring of blood glucose (SMBG) and (5) foot care. We kept the questions regarding medication adherence, physical activity, SMBG and foot care as they appeared on the SDSCA instrument, and we modified the questions regarding dietary behaviors to suit the local context. Firstly, Nepalese people tend to have 2 big meals per day, however, Nepalese health personnel generally recommend people with diabetes to have 5 small meals throughout the day. We decided to replace the question: "How many of the last 7 days have you followed a healthy eating plan?" with the question: "How many days (within the last 7 days) did you eat 5 or more small meals per day?". Furthermore, Nepalese people tend to eat one cup/bowl of vegetables per meal,

but fruit consumption varies, based on various factors. We; therefore, replaced the question: "On how many of the last 7 days did you eat five or more servings of fruits or vegetables?" with two separate questions. The first one being: "On how many of the last 7 days did you consume 1 cup/bowl of fruit per day?", and the second being: "On how many of the last 7 days did you consume 2 cups/bowls of vegetables per day?". In addition, an offering of sweets or mithai, which contain a high concentration of sugar, and are generally deep-fried in oil or ghee, is a common practice in Nepali culture during either festivals or on special occasions. Offers of sweets or mithai may come with undue pressure: it is considered polite to accept the offered sweets, while any refusal is considered rude. We thus added one additional question in the diet domain regarding the refusal of offered sweets/mithai within the past month.

#### **Measurement of perceived family support among people with type 2 diabetes**

We measured perceived family support using the DFBC." The DFBC consists of 21 items on a 5-point scale, reflecting the frequency of support provided by family members: (1) represents no support provided at all, (2) represents not more frequently than twice a month, (3) indicates more than twice a month, up to once a week, (4) represents more than once a week, up to several times a week, and (5) indicates at least once a day. The study instrument included five additional items, from the original DFBC, to suit the Nepali context of diabetes control, such as serving or requesting to have more vegetables than rice. White rice, which has a high glycemic index<sup>15</sup>, occupies the major portions of each meal in every household in Nepal. Healthcare practitioners generally recommend diabetic patients to have more vegetables than rice at each meal, so as to potentially lower the glycemic index of the meal. We also added items on serving, or requesting five

or more small meals per day; and whether family members did: “Not let you buy sweets” and: “Not let you eat offered sweets”, as well as whether family members “(Prepared) a separate meal for you during occasions”. This was done to adapt the study instrument to a Nepali context.

We checked the completed questionnaires for completeness prior to data entry, which was conducted using EpiData version 3.1. Univariate descriptive analysis was performed to present the distribution of basic characteristics of study participants, self-care activities, and family support. Exploratory factor analysis was performed to identify the underlying components of perceived family support. All data analyses were conducted using the R language and environment.

Regarding diabetes self-care behaviors, medication adherence was defined as: whether the respondent took their prescribed medicine/insulin on all seven days of the past week. In regards to physical activity, the original questionnaire measured physical activity using two items: frequency of physical activity related to work/household chores, and frequency of recreational physical activity. We combined these two items into one, and categorized participants into sedentary, moderately active and active. Sedentary patients included those who did not do any chores, nor performed any recreational physical activity during the past week. Moderately active patients referred to those who did some chores or performed recreational physical activity for fewer than 7 days in the past week, while active patients referred to participants who did chores or performed recreational physical activity for 7 days in the past week. In terms of dietary behaviors, we classified participants as having adherence to diet if; for the past 7 days, they had five or more small meals every day, consumed at least 2 bowls of vegetables every day and 1 or more bowls of fruit every day, consumed fatty food or red meat once or not at all, or within the past month, refused sweets when offered to them. We considered

participants who monitored blood glucose levels three or more times during the last seven days as having adequate adherence to SMBG, following the “guideline for self-monitoring of blood glucose in adults with diabetes”.<sup>16</sup> We defined foot care behavior adherence according to whether the participant checked his/her feet, and the inside of his/her shoes every day in the last seven days.<sup>17</sup>

We dichotomized responses to the family support items as: “Never engaged in the behavior” (response 1) vs. “Ever engaged in the behavior” (response 2–5), and presented the prevalence of family support behaviors using descriptive statistics. We also used this dichotomized scale in exploratory factor analysis (EFA). Of the 19 questions, on our modified DFBC, we excluded 2 items where the prevalence was over 90.0%; for example, items whose answers were too homogeneous to distinguish patterns (“eat at the same time as you” and “let you sleep rather than waking you up for medicine”). We created a scree plot to estimate the number of factors to identify, and performed EFA using orthogonal rotation. We retained questions where the factor loading exceeded 4.0 for the subsequent round of EFA, so as to create the most parsimonious factors. We found that 7 question items; for example, (How often do family members) “suggest things that help you to take your medicines/insulin on time?”, “help you to decide changes based on glucose results?”, “encourage you to participate in sports?”, “criticize you for not recording your blood glucose results?”, “eat other food that is not part of your diabetic diet?”, “not let you eat offered sweets?”, and “prepare a separate meal for you during special occasions?” were removed as all these variables that did not load on any factor with a loading higher than 0.40. To validate the identified family support factors, we derived the individual score for each factor, and assigned an arbitrary cut-off point at the 75<sup>th</sup> percentile of each factor, so as to classify the level of family support in the factor as: “High” (75<sup>th</sup> percentile or

higher) or: “Low” (below 75<sup>th</sup> percentile). These were then cross-tabulated with the prevalence of the dichotomized responses for each item of family support behavior.

This research was approved by the Institutional Review Board of Prince of Songkla University, Hat Yai, Thailand (REC. 62-204-19-9) and the Nepal Health Research Council (Reg. no. 588/2019).

## Results

We approached 428 eligible patients in the outpatient departments of TUTH, of which 411 consented to participate (96.0% response rate). More than half of the participants were female, and almost three-quarters were between the ages of 36 to 65 (Table 1). The most common ethnicity was Brahmin. Nearly all participants were married (93.7%), more than half earned less than 10,000 Nepalese Rupee [\$90 United States dollar (\$)] per month and 60.0% lived in a nuclear family. Three-quarters had had diabetes for more than 12 months, and nearly half were on two or more medications for diabetes. Most participants had at least one comorbidity; the most common being hypertension. Nearly all (98.3%) participants received a fasting blood sugar test, while testing for HbA1c was relatively uncommon (31.1%).

Among participants who were prescribed oral hypoglycemic agents and insulin, nearly all took medicines daily during the past 7 days (Table 2). Most participants had been physically active every day during the past 7 days. Adequate vegetable consumption was more common than adequate consumption of fruits, and having 5 or more small meals per day. More than half of the participants ate fatty food, or red meat only once, or had not during the last 7 days, and nearly three quarters had refused offered sweets within the past month. SMBG was not a common practice. Half of the participants had checked their feet every day within the last 7 days; whereas, less than a third had checked the inside of their shoes every day within the last seven days.

**Table 1** Basic characteristics and health profiles of study participants (n=411)

Characteristic	Frequency (%)
Gender: Female	234 (56.9)
Age group (years)	
21-35	47 (11.4)
36-50	119 (29.0)
51-65	177 (43.1)
66-80	64 (15.6)
81-95	4 (1.0)
Ethnicity	
Brahmin	125 (30.4)
Chhetri	67 (16.3)
Newar	60 (14.6)
Janajati	107 (26.0)
Dalit	17 (4.1)
Others	35 (8.5)
Marital status	
Never married	7 (1.7)
Married	385 (93.7)
Widowed	19 (4.6)
Current address	
Within Kathmandu	224 (54.5)
Outside Kathmandu	187 (45.5)
Education	
Illiterate (no formal education)	132 (32.1)
Literate (no formal education)	69 (16.8)
Primary school level (grades 1-5)	28 (6.8)
Lower secondary and secondary (grades 6-10)	32 (7.8)
School leaving certificate and higher secondary	100 (24.3)
University degree or higher	50 (12.2)
Income range (monthly; Nepali rupees)	
<10,000	216 (52.6)
10,000-20,000	96 (23.4)
>20,000-30,000	55 (13.4)
>30,000-40,000	27 (6.6)
>40,000	17 (4.1)
Family type	
Nuclear family	247 (60.1)
Joint/extended family	164 (39.9)
Duration of diabetes	
3-12 months	99 (24.1)
>12 months-5 years	167 (40.6)
>5-10 years	80 (19.5)
>10 years	65 (15.8)
Diabetes treatment	
Oral hypoglycemic agent only	284 (69.1)
Oral hypoglycemic agent and insulin	53 (12.9)
Insulin only	17 (4.1)

**Table 1** (continued)

Characteristic	Frequency (%)
None	55 (13.4)
Others	2 (0.5)
Number of drugs	
None	55 (13.4)
One	161 (39.2)
Two or more	187 (45.5)
Don't know	8 (1.9)
Comorbidities	
Hypertension	151 (36.7)
Chronic obstructive pulmonary disease	9 (2.2)
Cardiovascular diseases other than hypertension	58 (14.1)
Kidney disease	9 (2.2)
Cerebrovascular disease	10 (2.4)
Thyroid disease	34 (8.3)
Others <sup>1</sup>	33 (8.2)
Had one or more comorbidities	219 (53.3)
Fasting blood sugar (mg/dl) <sup>2</sup>	
Controlled ( $\leq$ 130)	175 (42.6)
Uncontrolled ( $>$ 130)	229 (55.7)
Not measured	7 (1.7)
Glycated hemoglobin level	
Controlled ( $<$ 7%)	45 (10.9)
Uncontrolled ( $\geq$ 7%)	83 (20.2)
Not measured	283 (68.9)

<sup>1</sup>dyslipidemia, psychiatric illness, hyperuricemia

<sup>2</sup>milligrams per deciliter

Table 3 shows a frequency list of family behaviors, as reported by the participants. The three most commonly reported family support behaviors were eating at the same time as the participants (98.1%), suggesting ways to help participants take medicine/insulin on time (89.5%), and helping participants to decide on changes based on glucose results (84.2%). The least commonly reported family support behaviors included letting the participants sleep, rather than rousing them to take their medicine (7.5%), criticizing the participants for not recording blood glucose results (29.2%) and criticizing the participants for not being physically active (33.3%).

**Table 2** Diabetes self-care practices of study participants (n=411 unless stated otherwise)

Self-care behaviors	Frequency (%)
Medication adherence	
Oral medicine (n=340 patients)	334 (98.2)
Insulin (n=70 patients)	70 (100.0)
Physical activity	
Sedentary*	23 (5.6)
Moderately active**	33 (8.0)
Active***	355 (86.4)
Dietary behavior	
Ate $\geq$ 5 small meals <sup>†</sup>	63 (15.3)
Ate $\geq$ 2 bowls of vegetables <sup>†</sup>	322 (78.3)
Ate $\geq$ 1 bowl of fruits <sup>†</sup>	186 (45.3)
Ate fatty food or red meat <sup>‡</sup>	228 (55.5)
Refused offered sweets within the past 1 month	287 (70.0)
Self-monitoring of blood glucose	
Monitored blood glucose level <sup>§</sup>	59 (14.4)
Foot care	
Checked feet <sup>†</sup>	210 (51.1)
Checked inside of shoes <sup>†</sup>	135 (32.8)

<sup>†</sup>Every day in the last 7 days. <sup>‡</sup>At most once in the last 7 days.

<sup>§</sup>At least three times in the last 7 days.

\*Sedentary: did not perform chores or recreational physical activity in the past 7 days

\*\*Moderately active: performed chores or recreational physical activity, for a total of  $<$ 7 days in the past 7 days

\*\*\*Active: performed chores or recreational physical activity every day in the past 7 days

Exploratory factor analyses yielded 3 factors of family support behaviors; as shown in Table 4. The behaviors that loaded onto the first factor included: nagging about testing blood sugar level, nagging about following the diet, arguing with the participant about self-care activities and preventing the participants from accepting offered sweets. As most of these activities were associated with control and criticism, we labeled this factor: "Authoritarian behavior". The behaviors that loaded onto the second factor included: praising the participants

**Table 3** Diabetes Family Behavior Checklist; as reported by study participants

Family behavior	“Ever” performed Frequency (%)
Ate at the same time as you	403 (98.1)
Suggested things that helped you to take medicines/insulin on time	368 (89.5)
Helped you to decide changes, based on your glucose results	346 (84.2)
Planned family activities that fit your diabetes self-care	329 (80.0)
Prepared a separate meal for you during special occasions	320 (77.9)
Bought things for you to carry in case of hypoglycemia	317 (77.1)
Served you/requested you to have more vegetables than rice	304 (74.0)
Served you/requested you to have 5 or more small meals	303 (73.7)
Nagged you about following a diet	300 (73.0)
Argued with you about diabetes self-care activities	296 (72.0)
Nagged about testing blood sugar level	295 (71.8)
Gave praise when you followed a diet	281 (68.4)
Prevented you from eating sweets offered to you	238 (57.9)
Congratulated you for adhering to your diabetes self-care	228 (55.5)
Ate other food that is not part of your diabetic diet	227 (55.2)
Encouraged you to participate in sports	175 (42.6)
Prevented you from buying sweets	162 (39.4)
Exercised with you	153 (37.2)
Criticized you for not being physically active	137 (33.3)
Criticized you for not recording your blood glucose results	120 (29.2)
Let you sleep rather than waking you to take your medicine	31 (7.5)

**Table 4** Results of factor analysis of perceived family support; according to diabetes patients (n=411)

Factor loadings	Factor 1 (Authoritarian)	Factor 2 (Supportive)	Factor 3 (Planning)
Praised you when you followed the diet	-0.121	0.565*	
Nagged you about testing blood sugar level	0.582*		
Criticized you for being physically inactive	0.359	0.512*	
Nagged you about following the diet	0.889*	-0.101	
Argued with you about diabetes self-care activities	0.738*		0.340
Planned family activities that fit your diabetes self-care routine	0.314	-0.112	0.762*
Congratulated you for adhering to your diabetes self-care routine		0.607*	
Exercised with you		0.426*	
Bought things for you to carry in case of hypoglycemia			0.514*
Served you/requested you to have more vegetables than rice		0.469*	
Prevented you from accepting offered sweets	0.404*		0.244
Served you/requested you to have 5 or more small meals		0.567*	
Percent variance explained	17.5%	14.3%	8.0%
Eigen values	2.1	1.7	1.0

\*Factor loading&gt;0.40

when following the diet, criticizing the participants for being physically inactive, congratulating the participants for adhering to self-care, exercising with the participants, serving or requesting the participants to eat more vegetables and to have smaller meals. As most of these behaviors were associated with showing support, we labeled this factor: "Supportive behavior". The behaviors that loaded onto the third factor included: planning family activities and buying things for the participants to carry in case of hypoglycemia. These behaviors were labelled as: "Planning behaviors". We validated the results of exploratory factor analyses using an arbitrary categorization of the behavioral levels, based on the factor analysis score of each factor, into high (above the 75<sup>th</sup> percentile) and low (75<sup>th</sup> percentile or lower) levels of "Authoritarian", "Supportive", and "Planning" behaviors. The prevalence of family support behaviors among those in the "high" group are shown in Table 5.

## Discussion

This study investigated sociodemographic and disease characteristics, self-care practices, and perceived

family support among people with type 2 diabetes receiving care at a tertiary center hospital in Kathmandu, Nepal. We used validated instruments, tailored to suit the context of the study setting, and found a high level of adherence to self-care behaviors in the domains of medication adherence and physical activity. However, adherence to dietary behaviors, self-monitoring of blood glucose, and foot care varied. Family support behaviors appeared to exist in three domains, but the prevalence of family support behaviors within each domain was very heterogeneous.

The most performed self-care practice in this study was medication adherence (>98.0%), followed by physical activities (>94.0%). This high percentage of medication compliance was also found in studies from Ethiopia, China and Malaysia.<sup>18-20</sup> The high level of compliance could be due to the influence of the study setting; wherein, there was a high level of trust in, and respect for, the attending clinicians.<sup>21</sup> Adherence to physical activity in our study was higher than in Malaysia, India and Ethiopia.<sup>18,20,22</sup> A possible reason is that we considered work/household related chores to be physical activity, unlike in other studies. Howbeit, this difference might also

**Table 5** Prevalence of reported family support among those with a high level (75<sup>th</sup> percentile ranking or higher) of authoritarian, supportive and planning domains of family support

Variables	Factor 1 (%) (Authoritarian)	Factor 2 (%) (Supportive)	Factor 3 (%) (Planning)
Praised you when you followed a diet	73.0	98.0	77.6
Nagged about testing blood sugar level	100.0	87.1	50.0
Criticized you for not being physically active	88.0	88.1	10.2
Nagged you about following a diet	100.0	76.2	48.0
Argued with you about diabetes self-care activities	100.0	73.3	63.3
Planned family activities that fit your diabetes self-care	88.0	77.2	100.0
Congratulated you for sticking in your diabetes self-care	69.0	97.0	76.5
Exercised with you	49.0	71.3	33.7
Bought things for you to carry in case of hypoglycemia	71.0	84.2	100.0
Served you/requested you to have more vegetables than rice	91.0	99.0	74.5
Prevented you eat offered sweets	86.0	62.4	56.1
Served you/requested you to have 5 or more small meals	81.0	100.0	95.9

be due to differences in lifestyles. Age could also have contributed to this difference, as more than four-fifths of our participants were less than 65 years of age, therefore they may have been able to perform daily physical activities.

Most of the participants had consumed two or more bowls of vegetables every day for the past week, whereas less than half had consumed one or more bowls of fruit. In Nepali cuisine, vegetables are a fundamental component of each meal, and families generally will make sure that each person has a cup of vegetables with each meal; regardless of their income. Fruit, on the other hand, is not considered as essential as vegetables, and will be consumed only by those who can afford it. Over half of all participants earned 10,000 Nepalese rupee (\$90 per month) or less, and this low level of income might explain the discrepancies between vegetable and fruit consumptions. Low consumption of fruit was also reported in studies from India.<sup>22,23</sup> The prevalence of fatty foods or red meat consumption in this study was higher than in previous studies<sup>22,23</sup>, which may be due to the timing of our study. This study was conducted in between two major festivals of Nepal (Dashain and Tihar); wherein, the primary dishes cooked at these festivals consist of goat's meat along with deep-fried food.

One-tenth of respondents adhered to SMBG, a result similar to findings reported in Ethiopia and Ghana.<sup>24,25</sup> The perceived, higher cost of self-monitoring may have been the cause of low adherence to SMBG.<sup>26</sup> The prevalence of checking one's own feet in this study was higher than in a study from Pakistan, but the prevalence of checking the inside of one's shoes was lower than the prevalence found in studies from Ethiopia and India.<sup>27-29</sup> Further, quantitative analyses and qualitative data are needed for a better understanding of the low level of foot care adherence found in our study.

In contrast to the two domains of family support behaviors presented in the original DFBC instrument, our

EFA identified 3 domains; for example, "Authoritarian", "Supportive", and "Planning". The "Authoritarian" behaviors were similar to the non-supportive behaviors' subscale in the DFBC, while the "Supportive" behaviors were like the DFBC's supportive behaviors subscale. The additional family support behaviors of "Serve/request to have more vegetables than rice" and "Serve/request to have five or more small meals" loaded on the Supportive behaviors domain, while "Not letting you eat offered sweets" loaded on the Authoritarian behaviors domain, showing consistency between the questions tailored to the Nepali context and the domains in the original DFBC. The similarities of these findings warrant further investigation.

There are several advantages in adapting data collection tools to suit the local context in the study of diabetes-related behaviors, and potential behavioral determinants. In regards to the healthcare system: the lack of economic resources in Nepal has implications on both access to medical care and the patient's lifestyle. Medical and lifestyle recommendations for patients in high-income countries may not be applicable, nor appropriate for patients in low-resource settings.<sup>30</sup> The value of food is also different in Nepal compared to high-income countries. Food is beyond nutrition; it is a means to develop, maintain and foster interpersonal relationships as well as bonds within and between both families and groups.<sup>31</sup> Hence, changes in dietary restrictions, or intake patterns can be misperceived as rejection or refusal of interpersonal relationships. Nepalese families function on a collective basis and give deference to their elders, and so these family values imply that family members may be reluctant to engage in conflicting behaviors, or to criticize diabetic patients who are often the more senior members of the family. Measuring food intake and family support in ways that are consistent with these unique values helps to create a better understanding of the distribution of behaviors, along with potential behavioral determinants

in a given population in manners that are relevant to actual health education and public health practices. To the best of our knowledge, this is the first study to measure diabetes self-care behaviors, and family support with standardized tools adapted to the Nepali context. However, several limitations should be taken into consideration. Firstly, we did not undergo multiple iterations for validation of the question items that were tailored to the Nepali context, which might have potentially undermined the validity of the findings. In addition, all responses in this study were self-reported. If patients believed that interviewers expected the participants to have a high level of adherence to self-care behaviors, social desirability could have influenced the responses, and the participants could have over-reported the prevalence of self-care behaviors. Lastly, this study was a hospital-based cross-sectional study, conducted in only one tertiary hospital in the capital city of Kathmandu. The findings of this study cannot be generalized to the context of rural Nepal, where most of the population lives and healthcare resources are more limited.

## Conclusion

We found variations in self-care activities and family support among people with type 2 diabetes attending outpatient services at a teaching hospital in Kathmandu, Nepal. Our study presented data from questions tailored to the local context, which may be helpful for future health promotion programs in the region as well as providing a methodological basis for future studies in other low and middle-income countries.

## Acknowledgement

We are grateful to all study participants, Tribhuvan University Teaching Hospital, and to Miss Lalla Maharjan and Miss Sanchita Karki for their help during the data collection process.

## Funding sources

The study was supported by a grant from the Thailand International Cooperation Agency.

## Conflict of interest

None

## References

1. International Diabetes Federation. IDF diabetes atlas [homepage on the Internet]. Brussels: IDF; 2017 [cited 2019 May 16]. Available from: <https://diabetesatlas.org/>
2. United Kingdom Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;352:837–53.
3. Ahmed Z, Yeasmeen F. Active family participation in diabetes self-care: a commentary. *Diabetes Manag* 2016;6:104–7.
4. World Health Organization. Adherence to long-term therapies: evidence for action. Geneva: WHO; 2003;p.198.
5. Rhee MK, Slocum W, Ziemer DC, Culler SD, Cook CB, El-Kebbi IM, et al. Patient adherence improves glycemic control. *Diabetes Educ* 2005;31:240–50.
6. D'Souza MS, Karkada SN, Parahoo K, Venkatesaperumal R, Achora S, Cayaban ARR. Self-efficacy and self-care behaviours among adults with type 2 diabetes. *Appl Nurs Res* 2017;36:25–32.
7. Fisher LW, Chesla CA, Bartz RJ, Gilliss C, Skaff MA, Sabogal F, et al. The family and type 2 diabetes: a framework for intervention. *Diabetes Educ* 1998;24:599–607.
8. Mayberry LS, Osborn CY. Family support, medication adherence, and glycemic control among adults with type 2 diabetes. *Diabetes Care* 2012;35:1239–45.
9. Tang TS, Brown MB, Funnell MM, Anderson RM. Social support, quality of life, and self-care behaviors among African Americans with type 2 diabetes. *Diabetes Educ* 2008;34: 266–76.
10. Scarton L, Velazquez I, O'Neal LJ, Iyerm S, Cannady T, Choate A, et al. Developing a culturally tailored multi-generational intervention to prevent and manage type 2 diabetes in American Indian families living in rural settings:

- Findings from a focus group study. *Res Nurs Health* 2019; 42:226–33.
11. Chesla CA, Chun KM, Kwan CML. Cultural and family challenges to managing type 2 diabetes in immigrant Chinese Americans. *Diabetes Care* 2009;32:1812–6.
  12. Sohal T, Sohal P, King-Shier KM, Khan NA. Barriers and facilitators for type-2 diabetes management in South Asians: a systematic review. *PLoS One* 2015;10. doi: 10.1371/journal.pone.0136202.
  13. Toobert DJ, Hampson SE, Glasgow RE. The summary of diabetes self-care activities measure: results from 7 studies and a revised scale. *Diabetes Care* 2000;23:943–50.
  14. Lewin AB, Geffken GR, Heidgerken AD, Duke DC, Novoa W, Williams LB, et al. The diabetes family behavior checklist: a psychometric evaluation. *J Clin Psychol Med Set* 2005;12: 315–22.
  15. Sun Q, Spiegelman D, van Dam RM, Holmes MD, Malik VS, Willett WC, et al. White rice, brown rice, and risk of type 2 diabetes in US men and women. *Arch Intern Med* 2010;170: 961–9.
  16. Wandsworth Clinical Commissioning Group. Self monitoring blood glucose guidelines [monograph on Internet]. London: Wandsworth Clinical Commissioning Group [cited 2020 Apr 26]. Available from: <https://www.wandsworthccg.nhs.uk/newsAndPublications/Publications/Diabetes/Self%20Monitoring%20Blood%20Glucose%20Guidelines.pdf>
  17. The Global Diabetes Community. Foot checks for diabetes [homepage on Internet]. Coventry: Diabetes Digital Media Ltd; 2019 [cited 2020 May 7]. Available from: <https://www.diabetes.co.uk/foot-care/foot-checks.html>
  18. Bongor Z, Shiferaw S, Tariku EZ. Adherence to diabetic self-care practices and its associated factors among patients with type 2 diabetes in Addis Ababa, Ethiopia. *Patient Pref Adher* 2018;12:963–70.
  19. Yin X, Savage C, Toobert D, Wei Pan, Whitmer K. Adaptation and testing of instruments to measure diabetes self-management in people with type 2 diabetes in Mainland China. *J Transcult Nurs* 2008;19:234–42.
  20. Gunggu A, Thon CC, Whye Lian C. Predictors of diabetes self-management among type 2 diabetes Patients. *J Diabetes Res* 2016;2016:9158943.
  21. Bhandari P, Kim M. Self-care behaviors of nepalese adults with type 2 diabetes: a mixed methods analysis. *Nurs Res* 2016;65:202–14.
  22. Rajasekharan D, Kulkarni V, Unnikrishnan B, Kumar N, Holla R, Thapar R. Self-care activities among patients with diabetes attending a tertiary care hospital in Mangalore Karnataka, India. *Ann Med Health Sci Res* 2015;5:59–64.
  23. Gopichandran V, Lyndon S, Angel MK, Manayalil BP, Blessy KR, Alex RG, et al. Diabetes self-care activities: a community-based survey in urban southern India. *Natl Med J India* 2012; 25:14–7.
  24. Berhe KK, Gebru HB, Kahsay HB, Kahsay AA, Kalayou. Assessment of self care management and its associated factors among type 2 diabetes patients in Mekelle Hospital and Ayder Referral Hospitals, Mekelle City, Tigray, Northern Ethiopia, 2012/13. *Global J Med Res* 2017;17:31–48.
  25. Mogre V, Abanga ZO, Tzelepis F, Johnson NA, Paul C. Adherence to and factors associated with self-care behaviours in type 2 diabetes patients in Ghana. *BMC Endocr Disord* 2017;17:20.
  26. Skeie S, Thue G, Nerhus K, Sandberg S. Instruments for self-monitoring of blood glucose: comparisons of testing quality achieved by patients and a technician. *Clin Chem* 2002;48:994–1003.
  27. Saeed N, Zafar J, Atta A. Frequency of patients with diabetes taking proper foot care according to international guidelines and its impact on their foot health. *J Pak Med Assoc* 2010;60: 732–5.
  28. Moreno Hernandez MI, Trilla Soler M, Espluga Capdevila A, Mengual Miralles N, Bundo Vidiella M, et al. Self care and risk factors of diabetic foot in patients with type II diabetes mellitus. *Aten Primaria* 1997;20:185–90.
  29. Uma MR, Sowiya KR, Kavin S. Self-care practices among type II diabetics attending primary health centre, Thiruvallur district, Tamil Nadu. *Int J Community Med Public Health* 2017;4:2745–9.
  30. Dagenais GR, Leong DP, Rangarajan S, Lanans F, Lopez-Jaramillo P, Gupta R, et al. Variations in common diseases, hospital admissions, and deaths in middle-aged adults in 21 countries from five continents (PURE): a prospective cohort study. *Lancet* 2020;395:785–94.
  31. Prakash J, Waisundara V, Prakash V. Nutritional and health aspects of food in South Asian Countries. Cambridge: Academic Press; 2020;p.365.